

DTIC FILE COPY

①


AD-A227 123



# JOINT U.S. NAVY/U.S. AIR FORCE CLIMATIC STUDY OF THE UPPER ATMOSPHERE


## VOLUME 2 - FEBRUARY

SEPTEMBER, 1989



PREPARED BY  
NAVAL OCEANOGRAPHY COMMAND DETACHMENT  
ASHEVILLE, N.C.

PREPARED UNDER THE AUTHORITY OF  
COMMANDER, NAVAL OCEANOGRAPHY COMMAND  
STENNIS SPACE CENTER, MS 39529-5000



DTIC  
ELECTED  
OCTO 4 1990  
S E D



08501.P0155500



UNCLASSIFIED

SECURITY CLASSIFICATION OF THIS PAGE

REPORT DOCUMENTATION PAGE					
1a. REPORT SECURITY CLASSIFICATION Unclassified		1b. RESTRICTIVE MARKINGS			
2a. SECURITY CLASSIFICATION AUTHORITY		3. DISTRIBUTION/AVAILABILITY OF REPORT Public Release/Distribution Unlimited			
2b. DECLASSIFICATION/DOWNGRADING SCHEDULE					
4 PERFORMING ORGANIZATION REPORT NUMBER(S)		5. MONITORING ORGANIZATION REPORT NUMBER(S) NAVAIR 50-1C-2 S/N 0850-LP-015-5500, AWS/TR-89/002			
6a. NAME OF PERFORMING ORGANIZATION National Climatic Data Center Global Analysis Branch		6b OFFICE SYMBOL (If applicable) E/CC22	7a. NAME OF MONITORING ORGANIZATION Naval Oceanography Command Detachment Asheville		
6c. ADDRESS (City, State, and ZIP Code) Federal Building Asheville, NC 28801-2696		7b. ADDRESS (City, State, and ZIP Code) Federal Building Asheville, NC 28801-2696			
8a. NAME OF FUNDING/SPONSORING ORGANIZATION <u>Commander, Naval Oceanography Command</u> <u>Headquarters, Air Weather Service</u>		9. PROCUREMENT INSTRUMENT IDENTIFICATION NUMBER			
8c. ADDRESS (City, State, and ZIP Code) <u>Stennis Space Center, MS 39529-5000</u> Scott AFB, IL 62225-5008		10. SOURCE OF FUNDING NUMBERS			
		PROGRAM ELEMENT NO.	PROJECT NO.	TASK NO.	WORK UNIT ACCESSION NO.
11. TITLE (Include Security Classification) Joint U.S. Navy/U.S. Air Force Climatic Study of the Upper Atmosphere Volume 2-February					
12. PERSONAL AUTHOR(S) NCDC - Michael J. Changery, Claude N. Williams NAVOCEANCOMDET - Michael L. Dickenson, Brian L. Wallace					
13a. TYPE OF REPORT Final	13b. TIME COVERED FROM _____ TO _____	14 DATE OF REPORT (Year, Month, Day) September 1989		15. PAGE COUNT 236	
16. SUPPLEMENTARY NOTATION					
17. COSATI CODES		18. SUBJECT TERMS (Continue on reverse if necessary and identify by block number)			
FIELD	GROUP				
19. ABSTRACT (Continue on reverse if necessary and identify by block number)  This study of the upper atmosphere is based on 1980-85 twice daily gridded analysis produced by the European Centre for Medium Range Weather Forecasts. Included are global analyses of (1) Mean Temperature/Standard Deviation, (2) Mean Geopotential Height/Standard Deviation, (3) Mean Density/Standard Deviation, (4) Height and Vector Standard Deviation. All for 13 pressure levels - 1000, 850, 700, 500, 400, 300, 250, 200, 150, 100, 70, 50, 30 mb. In addition, analyses of (5) Mean Dew Point/Standard Deviation - levels 1000 through 300 mb, (6) jet stream (mean scalar speed) - levels 500 through 30 mb. Also included are global 5 degree grid point wind roses for the 13 pressure levels.					
20. DISTRIBUTION/AVAILABILITY OF ABSTRACT <input type="checkbox"/> UNCLASSIFIED/UNLIMITED <input checked="" type="checkbox"/> SAME AS RPT <input type="checkbox"/> DTIC USERS			21. ABSTRACT SECURITY CLASSIFICATION Unclassified		
22a. NAME OF RESPONSIBLE INDIVIDUAL Brian L. Wallace			22b. TELEPHONE (Include Area Code) (704) 252-7865	22c. OFFICE SYMBOL	

DD FORM 1473, 84 MAR

83 APR edition may be used until exhausted.

All other editions are obsolete.

SECURITY CLASSIFICATION OF THIS PAGE

UNCLASSIFIED

## TABLE OF CONTENTS

	PAGE
<b>INTRODUCTION . . . . .</b>	<b>iv</b>
<b>REFERENCES . . . . .</b>	<b>vii</b>
<b>ELEMENTS</b>	
<b>PRESSURE-HEIGHT. . . . .</b>	<b>1-27</b>
<b>WIND ROSE. . . . .</b>	<b>29-107</b>
<b>JET STREAM . . . . .</b>	<b>109-129</b>
<b>TEMPERATURE. . . . .</b>	<b>131-157</b>
<b>DEW POINT. . . . .</b>	<b>159-171</b>
<b>DENSITY. . . . .</b>	<b>173-199</b>
<b>HEIGHT/WIND STANDARD DEVIATION . . . . .</b>	<b>201-227</b>

Accession For	
NTIS GRA&I	<input checked="" type="checkbox"/>
DTIC TAB	<input type="checkbox"/>
Unannounced	<input type="checkbox"/>
Justification	
By _____	
Distribution/ _____	
Availability Codes	
Dist	Avail and/or Special
A-1	

The Joint U.S. Navy/U.S. Air Force Climatic Study of the Upper Atmosphere was prepared by the Officer in Charge, Naval Oceanography Command Detachment, Asheville, North Carolina under the authority of Commander, Naval Oceanography Command. Additional funding was provided by the Air Weather Service as a result of Tri-Services Climatology initiatives. The work was performed at the National Climatic Data Center (NCDC). Specific acknowledgement of the NCDC staff is made to Mr. M.J. Changery, project leader; Mr. C.N. Williams, Jr. for data processing and software development; and Messrs. M.G. Burgin and D.A. McKittrick for drafting skills. Special acknowledgement is made to the European Centre for Medium-range Weather Forecasts for providing the basic gridded analyses.

## INTRODUCTION

During the past decade, improvements in the collection and assimilation of data required for more accurate representations of the atmosphere have resulted in data sets useful for developing a more definitive climatology of the global atmosphere. Such a climatology has uses in aircraft operations and planning, indirect assessments of atmospheric transport as well as a standard state from which atmospheric anomalies can be analyzed.

Prior climatologies, U.S. Navy (1959), U.S. Navy (1966), Naval Weather Service Command (1969), and Naval Weather Service Command (1970), were produced from individual station data with varying periods of record, and the resulting summarized data were analyzed. A serious deficiency was the lack of reporting locations in the major ocean basins. Analyses over the oceans were derived by extrapolating from known analyses over coastal regions as well as the few island or ocean vessels available. An additional complication was the manually intensive effort required to ensure horizontal and vertical consistency of the data.

With the advent, in the 1970s, of more powerful computers and data collection and assimilation systems, the initial analyses used for input into forecast models had a three-fold advantage over the station analyses utilized in the prior climatologies. First, the data assimilation system utilized a greater variety of information for production of an analysis. The normal array of land-based upper air reporting stations was supplemented by ship-based reporting stations, cloud reports, pilot reports and, most importantly, satellite-derived temperature, moisture and wind data. Consequent analyses more accurately represented the state of the atmosphere at a given observation time. Second, the assimilation system quality-controlled all incoming data and ensured the horizontal and vertical consistency of the resulting analyses. Finally, through the computer-based system, global data were available and archived in grid-point form.

A number of analysis sets produced by various national and international meteorological services were investigated. It is recognized that improvements to the data assimilation and analysis systems occurred within any analysis set produced, and that current analyses more accurately reflect the atmosphere's state than do the earlier analyses. It is also recognized that specific parameter or geographic-based deficiencies exist in all analysis sets. However, the intent of this upper-air climatology effort is the production of analyses to serve the needs of the operational meteorologist. A climatology derived from global analyses achieves this goal. Based on known capabilities and technical reviews of the various systems, as well as recommendations from the professional numerical modeling community, the analyses produced by the European Centre for Medium-range Forecasts were selected for processing.

## ECMWF DATA

The European Centre for Medium-range Weather Forecasts (ECMWF) is an international organization established in 1973 and supported by 17 member states. It is responsible for providing global forecasts to the European community. Their data assimilation system consists of multivariate optimal interpolation analysis allowing the incorporation of a variety of observations with differing error characteristics and spatial distributions. A relatively comprehensive coverage of global data is ensured through the data collection schedule. A unique feature of the ECMWF system is the method of grid point analysis. Rather than analyzing individual grid points, varying sized boxes (depending on data density) are created containing groups of grid points. Grid point analysis uses data from within the box as well as adjacent boxes, thereby assuring a consistent analysis between all the grid points.

The system also includes internal quality control which examines the climatological reasonability of incoming data as well as the internal consistency of the data.

In addition, the system utilizes a model initialization process which ensures that harmful gravity waves, caused by imbalances in the analysis, with the potential to create problems in subsequent forecast fields, are suppressed. Through the initialization process, the atmosphere's mass and wind fields are adjusted so that only a portion of the gravity wave balanced by dynamic and physical processes is retained. Further information on the ECMWF system is available in Lorenc (1981), Shaw, et al. (1984), Lonnberg, et al. (1986), and ECMWF (1988).

The resulting initialized analyses are vertically interpolated to these 13 standard pressure levels: 1000, 850, 700, 500, 400, 300, 250, 200, 150, 100, 70, 50, and 30 mb, and include the geopotential height, temperature, and wind for all levels with moisture included for the 1000 through 300 mb levels.

Six years (1980-1985) of individual analysis were obtained from ECMWF on a 2.5° global grid. Although the analyses were permanently archived as spherical harmonic coefficients, ECMWF reconstituted the analyses for use in the data processing. Synoptic analyses at six-hour intervals were received for the six-year period, but only the 00 and 12Z analyses were re-sorted into a grid point sort. Given the quality control performed by ECMWF on collected data and the requirements for horizontal and vertical data consistency imposed by the assimilation system, minimal quality control was performed prior to summarization. Primary quality control was limited to comparison of level data against known/estimated climatological extremes.

The summarized grid point data were objectively analyzed, machine-contoured by parameter and level on polar stereographic (0°-90°N and S) and cylindrical equidistant (0°-60°N and S) projections with resulting contours machine-labeled. In addition, individual wind observations were consolidated into eight 45° segments centered on directions north, northeast, .... through northwest for display as wind roses on a series of cylindrical equidistant projections.

Since the ECMWF analyses were archived as spectral harmonic coefficients, the grid point reconstitution process provides data for all global 2.5° grid points. This naturally includes (for the 1000 through 700 mb levels) selected grid points at which the land elevations exceed the height of the pressure surface. For these grid points, a blanking program was used to eliminate both contours and grid point wind roses.

## ANALYSES

### 1. Pressure-Height

Grid point geopotential height values (in dekameters) are summarized by month for 13 levels from 1000 mb to 30 mb with solid and dashed contours of mean values presented on pressure height charts. Standard deviation of height is calculated from the individual daily values with contours presented on a separate chart series including the standard deviation of vector mean wind. Local points of highest and lowest pressure are designated with H's and L's on the analyzed charts. Not all pressure centers are enclosed by closed contours. Vector mean wind in 5-knot increments are calculated for selected grid points considered adequate to depict flow for the hemisphere with wind shaft orientation related to specific latitude/longitude lines. Vector mean winds less than 2.5 knots are depicted as a shaft with no barbs. Contours of mean geopotential height and vector mean wind barbs are presented for the northern/southern hemispheres on polar stereographic projection and for 0° to 60° north and south on cylindrical equidistant projections with blanking for appropriate high elevation land areas on the 1000 through 700 mb charts.

### 2. Wind Roses

Wind roses for 10° grid points from 5° to 85° north and south are presented by month for all levels from 1000 mb to 30 mb. Each hemisphere is divided into three longitudinal zones: 60°W to 60°E, 60°E to 180°E, and 180°W to 60°W. Each rose presents:

- a) Scalar mean speed
- b) Percent frequency of occurrence from each of 8 cardinal point wind directions proportional to shaft length with dots on the shafts representing 5 percentile intervals.
- c) Mean speed for each of the 8 cardinal wind directions rounded to the nearest 5 knots.

Roses for grid points on the 1000 mb through 700 mb level charts are blanked whenever the land elevation exceeds the mean geopotential height of the specified level.

### 3. Temperature

Grid point temperature data (in °C) are summarized by month for 13 levels from 1000 mb to 30 mb with solid and dashed contours of mean values presented on pressure height charts. Temperature standard deviation derived from the individual observations are shown on the same charts with dotted contours. Contours are presented for both the northern and southern hemispheres on a polar stereographic projection and for the zone from 0° to 60° north and south on cylindrical equidistant projections with blanking for appropriate high elevation land areas on the 1000 through 700 mb charts.

### 4. Dew Point

Grid point moisture data were received as mixing ratios for the period through April 19, 1982 and as relative humidity thereafter for the 1000 through 300 mb levels. All moisture data were converted to dew point values. These are summarized by month with solid and dashed contours of mean values presented on pressure height charts. Dew point standard deviation derived from the individual observations are shown on the same charts with dotted contours. Contours are presented for both the northern and southern hemispheres on a polar stereographic projection and for the zone from 0° to 60° north and south on cylindrical equidistant projections with blanking for appropriate high elevation land areas on the 1000 through 700 mb charts.

## 5. Density

Grid point density data were computed from the daily values of temperature and pressure from the equation of state in the form

$$\rho = \frac{P}{RT}$$

where  $\rho$  is the density, P is the pressure, T is the temperature, and R is the gas constant. Density was computed for moist air through 300 mb and for dry air from 250 mb to 30 mb. Density data (in Kg/m<sup>3</sup>) are summarized by month for all 13 levels with solid and dashed contours of mean values presented on pressure height charts. Density standard deviation derived from individual observations are shown on the same charts with dotted contours. Contours are presented for both the northern and southern hemispheres on a polar stereographic projection and for the zone from 0° to 60° north and south on cylindrical equidistant projections with blanking for appropriate high elevation land areas on the 1000 through 700 mb charts.

## 6. Standard Deviation of Height and Vector Mean Wind

Standard deviation of the height and vector mean wind data presented on the pressure height charts are presented on monthly charts for the 1000 through 30 mb levels. Height standard deviations (in dekameters) are presented as solid contours and vector wind standard deviations (in knots) as dashed contours. Contours are presented for both the northern and southern hemispheres on a polar stereographic projection and for the zone from 0° to 60° north and south on cylindrical equidistant projections with blanking for appropriate high elevation land areas on the 1000 through 700 mb charts.

## 7. Jet Stream

Grid point scalar mean wind speed (in knots), as presented by the value in the center of the wind rose octagons, are summarized by month and analyzed for 500 through 30 mb. All speeds exceeding 50 knots are shaded with shading intensity increasing by 25-knot increments. Contours are presented for both the northern and southern hemispheres on a polar stereographic projection and for the zone from 0° to 60° north and south on cylindrical equidistant projections.

## DATA AVAILABILITY

Monthly summarized grid point data for the period of record for all levels from 1000 through 30 mb have been retained on magnetic tape. Data available, per level, include:

- Number of observations
- Mean zonal wind component and standard deviation
- Mean meridional wind component and standard deviation
- Vector mean wind and standard deviation
- Mean temperature and standard deviation
- Mean dew point (through 300 mb) and standard deviation
- Mean geopotential height and standard deviation
- Mean density and standard deviation
- Mean scalar wind speed and percentage of observations for each designated direction

Similarly summarized data for each half-month of the 1980-85 period are also available on magnetic tape. Summaries can be provided on magnetic media or in listing form by the National Climatic Data Center.

## REFERENCES


- ECMWF, 1988: User guide to ECMWF products.
- Lorenc, A.C., 1981: A global three-dimensional multivariate statistical interpretation scheme. Monthly Weather Review, **109**, 701-721.
- Lonnberg, P., J. Pailleux, and A. Hollingsworth, 1986: The new analyses system. ECMWF Technical Memorandum No. 125.
- Naval Weather Service Command, 1969: Climate of the Upper Air - Southern Hemisphere, VOL I, Temperature, Dewpoint and Heights at Selected Pressure Levels, NAVAIR 50-1C-55.
- Naval Weather Service Command, 1970: Selected Level Heights, Temperatures and Dewpoints for the Northern Hemisphere, NAVAIR 50-1C-52.
- Shaw, D.B., P. Lonnberg, and A. Hollingsworth, 1984: The 1984 revision of the ECMWF Analysis System. ECMWF Technical Memorandum, No. 92.
- U.S. Navy, 1959: Upper Wind Statistics Charts of the Northern Hemisphere, VOL I-III, NAVAIR 50-1C-535.
- U.S. Navy, 1966: Components of the 1000 mb Winds of the Northern Hemisphere, NAVAIR 50-1C-51.

PRESSURE - HEIGHT  
(13 LEVELS, 1000 TO 30 MB)

- Contours of mean height (solid and dashed lines) in geopotential dekameters; example: 580 is 5800 geopotential meters; solids labeled, dashed intermediates unlabeled
- Height labeled interval:

6 dekameters (60 meters) - 1000 MB to 400 MB  
12 dekameters (120 meters) - 300 MB to 200 MB  
8 dekameters (80 meters) - 150 MB to 30 MB
- Vector mean wind in knots
- Contours blanked for geographic areas with elevations exceeding specified geopotential heights

ELEVATION SCALE





Mean Geopotential Height (dkm)

Vector Mean Wind (kt)

February



1000 Mb

Upper Air Climatology  
Northern Hemisphere



Upper Air Climatology  
Southern Hemisphere

Mean Geopotential Height (dkm)  
Vector Mean Wind (kt)  
February  
1000 Mb



Mean Geopotential Height (dkm)


Vector Mean Wind (kt)

February

850 Mb

Upper Air Climatology

Northern Hemisphere



Upper Air Climatology


Southern Hemisphere

Mean Geopotential Height (dm)

Vector Mean Wind (ft)

February

850 MB




### Mean Geopotential Height (dkm)

### Vector Mean Wind (kt)

February

700 MB

## Upper Air Climatology Northern Hemisphere





Upper Air Climatology  
Southern Hemisphere

Mean Geopotential Height (dkm)

Vector Mean Wind (kt)

February

700 hPa



Mean Geopotential Height (dkm)


Vector Mean Wind (kt)

February

500 Mb

Upper Air Climatology

Northern Hemisphere




Upper Air Climatology  
Southern Hemisphere

Mean Geopotential Height (dkm)

Vector Mean Wind (kt)

February

500 Mb



Mean Geopotential Height (dkm)


Vector Mean Wind (kt)

February

400 Mb


Upper Air Climatology

Northern Hemisphere



Upper Air Climatology  
Southern Hemisphere

Mean Geopotential Height (dkm)  
Vector Mean Wind (kt)  
February  
400 Mb



Mean Geopotential Height (dkm)


Vector Mean Wind (kt)

February

300 Mb


Upper Air Climatology

Northern Hemisphere



Upper Air Climatology  
Southern Hemisphere

Mean Geopotential Height (dkm)  
Vector Mean Wind (kt)  
February  
300 Mb



Mean Geopotential Height (dkm)



Vector Mean Wind (kt)

February

250 Mb

Upper Air Climatology

Northern Hemisphere





Upper Air Climatology  
Southern Hemisphere

Mean Geopotential Height (dkm)

Vector Mean Wind (kt)

February

250 Mb



Mean Geopotential Height (dkm)


Vector Mean Wind (kt)

February

200 Mb


Upper Air Climatology

Northern Hemisphere



Upper Air Climatology  
Southern Hemisphere

Mean Geopotential Height (dkm)  
Vector Mean Wind (kt)  
February  
200 Mb



Mean Geopotential Height (dkm)



Vector Wind (km)

February

100 hPa

Upper Air Climatology

Northern Hemisphere



Upper Air Climatology


Northern Hemisphere

Mean Geopotential Height (dkm)

Vector Mean Wind (kt)

February

150 Mb



### Mean Geopotential Height (dkm)



### Vector Mean Wind (kt)

February

100 MB

## Upper Air Climatology

### Northern Hemisphere




Upper Air Climatology  
Southern Hemisphere

Mean Geopotential Height (dkm)

Vector Mean Wind (kt)

February

100 Mb



Mean Geopotential Height (dkm)



Vector Mean Wind (kt)

February

70 Mb

Upper Air Climatology

Northern Hemisphere





## Upper Air Climatology Southern Hemisphere

### Mean Geopotential Height (dkm)

### Vector Mean Wind (kt)

February

70 Мб




Mean Geopotential Height (dkm)

Vector Mean Wind (kt)

February

50 Mb

Upper Air Climatology  
Northern Hemisphere





## Upper Air Climatology Southern Hemisphere

### Mean Geopotential Height (dkm)

### Vector Mean Wind (kt)

February

50 MB






## Upper and Lower Cervical Hemispheres

### Messung der Kapazität des Zellkerns (cm<sup>3</sup>)

1922-1923 (part) (2)


2000

卷之三





**WIND ROSES**  
**(13 LEVELS, 1000 TO 30 MB)**


- Wind roses at 10 degree latitude/longitude grid points
- Directional mean wind speed in 5 knot increments
- Frequency proportional to barb length with individual dots representing 5% increments. Values greater than 30% are plotted directly on the barb.
- Roses blanked at grid points with elevations exceeding specified geopotential heights.
- Sample rose explanation:




ELEVATION SCALE











Upper Air Climatology  
Southern Hemisphere

Climatic Data  
1950-1959

Probability  
1000 Metres






Upper Air Climatology  
Southern Hemisphere

12000 FT. ELEV.  
WINDS AT 1000 MB


February  
1000 MB



WORLD AIR CHART  
Northern Hemisphere

CHARTS  
NORTH HEMISPHERE


February  
1950



Upper Air Climatology  
Southern Hemisphere

5000 500 500  
Wind Wind Wind


February  
850 Mb



ପ୍ରକାଶକ  
ବ୍ୟାପକ  
ମେଲା

850 MHz


## **Upper Air Climatology Northern Hemisphere**



Upper Air Climatology  
Southern Hemisphere

SUITE NO. 6273  
WINDS 10.5232


February  
850 MB




Geographic  
Distribution  
of Species

Upper Air Climatology  
Northern Hemisphere

Geography  
250 m







Upper Air Climatology  
Northern Hemisphere

Figure 502  
Vertical Profile

Reference  
700 mb






February  
700 MB

50°E TR 1300Z  
Wind Roses


Upper Air Climatology  
Northern Hemisphere



Upper Air Climatology  
Southern Hemisphere

50E TO 180E  
Wind Rose


February  
700 Mb



四百九

ବ୍ୟାକ ପରିଚୟ


## Upper Air Climatology Northern Hemisphere




Upper Air Climatology  
Southern Hemisphere

Geophysical Society  
Washington  
1963

February  
700 MB






Upper Air Climatology  
Southern Hemisphere

Climatic Data  
1950-1953


Probability  
500 M.B.



500E 70° 1800E  
Wind roses

Frequency  
500 Miles


Upper Air Climatology  
Northern Hemisphere



Upper Air Climatology  
Southern Hemisphere

500 TO 1000  
Wind Rose


February  
500 MS



POLYMER  
500 M.E.

HOOTY TO SOOTY  
WIND ROSES


Upper Air Climatology  
Northern Hemisphere



Upper Air Climatology  
Southern Hemisphere

180°W TO 60°W  
Wind Rose


February  
500 Mb



## Upper Air Climatology Northern Hemisphere

డెస్కో డెవలా


400 M



Upper Air Climatology  
Southern Hemisphere

50°W 50°S  
Wind Rose


February  
400 MB



REGULARITY  
400 METS

STORM LOGS  
WIND ROSES


Upper Air Climatology  
Northern Hemisphere



## Upper Air Climatology Southern Hemisphere


SACRED PERIOD.

Geodiversity  
400 M€



WIND ROSES  
1000 METERS


Upper Air Climatology  
Northern Hemisphere



Upper Air Climatology  
Southern Hemisphere

1000 1500 2000  
mb mb mb


February  
1950




February  
300 Mil

Snow TDO 50E  
Wind Rose

Upper Air Climatology  
Northern Hemisphere






February  
300 MB

卷之三


## Upper Air Climatology Southern Hemisphere



Upper Air Climatology  
Southern Hemisphere

SOUTH HEMISPHERE  
UPPER AIR CLIMATOLOGY


February  
1952



卷之三


卷之三

## Upper Air Climatology Northern Hemisphere



## Upper Air Climatology Southern Hemisphere

卷之三



Upper Air Climatology  
Northern Hemisphere


July 1952  
Vol. 1, No. 2

Geophysical  
Books



Upper Air Climatology  
Southern Hemisphere


February 1953  
250 mb




Snow and Rain  
Water Precip.

Upper Air Climatology  
Northern Hemisphere

Geography  
Sect. No. 2









Upper Air Climatology  
Southern Hemisphere


February 1950  
250 mb



Geographic  
Upper Air Climatology  
Northern Hemisphere  
General Processes

Upper Air Climatology  
Northern Hemisphere






February  
2000


Upper Air Climatology  
Southern Hemisphere

અનુભૂતિ

卷之三

## Upper Air Climatology Northern Hemisphere






Upper Air Climatology  
Southern Hemisphere

Fig. 10.13  
Upper air climatology  
Southern Hemisphere

FEBRUARY  
1965

Upper Air Chemistry  
Northern Hemisphere


Upper Air Chemistry  
Northern Hemisphere




Upper Air Climatology  
Southern Hemisphere

20°S - 60°S  
1960-1970  
100 mb


Hadley Cell  
40°S MSL





Upper Air Climatology  
Southern Hemisphere


February 1950  
150 mb



February  
150 MN

Upper Air Climatology  
Wind Processes


Upper Air Climatology  
Northern Hemisphere



Upper Air Climatology  
Southern Hemisphere

500 MB ISONE  
WIND REGIONS


POLARITY  
150 MB



FEBRUARY  
150 M.

JANUARY TO JUNE  
WINDS & PRESS.


Upper Air Climatology  
Northern Hemisphere



Upper Air Climatology  
Southern Hemisphere

30°S 35°S  
40°S 45°S


February  
1950




FEBRUARY  
1000 MB

2000 2500 3000  
WINDS AND PRESSURE

Upper Air Climatology  
Northern Hemisphere





February  
100 hPa  
100 MB


Southern Hemisphere  
Upper Air Climatology

Southern Hemisphere

February  
100 Mb

Feb 1958 1800  
Wind Rosettes


Upper Air Climatology  
Northern Hemisphere




Upper Air Climatology  
Southern Hemisphere


Fig. 1303  
100 mb

February  
100 mb



## Upper Air Climatology Northern Hemisphere






## Upper Air Climatology Northern Hemisphere

卷之三


卷之三



Upper Air Climatology  
Southern Hemisphere

500 mb 500 mb  
February 1950


February  
70 Mb



## Upper Air Climatology Northern Hemisphere

50IE TO 180E  
Wined Rosess


40 مکالمہ



Upper Air Climatology  
Southern Hemisphere

50°E TO 180°E  
Wind Roses


February  
70 Mb



Upper Air Climatology  
Northern Hemisphere

12000 15000 6000  
Wind Rose


Scattering  
70 M.



Upper Air Climatology  
Southern Hemisphere

1950-51 Season  
70 Mb


February  
70 Mb



50° N 50° S  
Wind roses

Frequency  
50 MHz


Upper Air Climatology  
Northern Hemisphere





Upper Air Climatology  
Southern Hemisphere

50 mb 40° S  
WINDS

February  
50 mb







February  
50 Mb


50IE TO 180IE


## Upper Air Climatology Southern Hemisphere

30°W TO 60°W  
Wind roses

February  
50 N.E.

Upper Air Climatology  
Northern Hemisphere







February  
50 Mb

مکتبہ ملک

## Upper Air Climatology Southern Hemisphere





Tetbury  
30 Mb


卷之三

# Upper Air Climatology Southern Hemisphere

Upper Air Climatology  
Northern Hemisphere

FIGURE 1802  
Whirl Rose


February  
30 Mid




Upper Air Climatology  
Southern Hemisphere

500 mb 1000  
WINDS 30 SEC

February  
30 Mb






Upper Air Climatology  
Southern Hemisphere

1950-51  
1951-52


February  
30 M<sup>o</sup>



**JET STREAM  
(10 LEVELS, 500 TO 30 MB)**

- Contours of mean scalar wind speed in knots
- Minimum mean scalar speed: 50 knots
- Contour interval of mean scalar speed: 25 knots

**ELEVATION SCALE**




Jet Stream  
50kt + 25kt inc  
February  
500 MB

Upper Air Climatology  
Northern Hemisphere




Upper Air Climatology  
Southern Hemisphere

Jet Stream  
50kt + 25kt inc  
Relatively  
500 Mb




Jet Stream  
50kt + 25kt inc  
February  
400 MB

Upper Air Climatology  
Northern Hemisphere





Upper Air Climatology  
Southern Hemisphere

Jet Stream  
50kt + 25kt inc  
February  
400 Mb




Jet Stream  
50kt + 25kt inc  
February  
300 Mb

Upper Air Climatology  
Northern Hemisphere





Upper Air Climatology  
Southern Hemisphere

Jet Stream  
50kt + 25kt inc  
February  
300 Mb





Jet Stream  
50kt + 25kt inc  
February  
250 Mb

Upper Air Climatology  
Northern Hemisphere



Upper Air Climatology  
Southern Hemisphere

Jet Stream  
50kt + 25kt inc  
February  
250 MB



Jet Stream



50kt + 25kt inc

February

200 Mb

Upper Air Climatology

Northern Hemisphere



Upper Air Climatology


Southern Hemisphere

Jet Stream

50kt + 25kt inc


February

200 Mb



Jet Stream  
50kt + 25kt inc  
February  
150 MB

Upper Air Climatology  
Northern Hemisphere



Upper Air Climatology



Southern Hemisphere

Jet Stream

50kt + 25kt inc


February

150 Mb



Jet Stream  
50kt + 25kt inc  
February  
100 Mb

Upper Air Climatology  
Northern Hemisphere



Upper Air Climatology  
Southern Hemisphere

Jet Stream  
50kt + 25kt inc  
February  
100 Mb


Jet Stream  
50kt + 25kt inc  
February  
70 Mb

Upper Air Climatology  
Northern Hemisphere

50° W 0° 50° E 90° E


Upper Air Climatology  
Southern Hemisphere

Jet Stream  
50kt + 25kt inc  
February  
70 Mb



Jet Stream  
50kt + 25kt inc  
February  
50 Mb

Upper Air Climatology  
Northern Hemisphere




Upper Air Climatology  
Southern Hemisphere

Jet Stream  
50kt + 25kt inc  
February  
50 Mb

50 Wind < 50 Kt


Jet Stream  
50kt + 25kt inc  
February  
30 Mb

Upper Air Climatology  
Northern Hemisphere



Upper Air Climatology  
Southern Hemisphere


Jet Stream  
50kt + 25kt inc  
February  
30 MLS



**TEMPERATURE**  
(13 LEVELS, 1000 TO 30 MB)

- Contours of mean temperature (solid and dashed lines) in °C; solids labeled, dashed intermediates unlabeled
- Temperature labeled interval: 5°C
- Contours of standard deviation of temperature (dotted lines) in °C
- Standard deviation of temperature labeled interval: 2.5°C
- Contours blanked for geographic areas with elevations exceeding specified geopotential heights

**ELEVATION SCALE**



Mean Temperature (°C)



Std Dev (Dotted)

February

1000 mb


Upper Air Climatology

Northern Hemisphere



Upper Air Climatology  
Southern Hemisphere

Mean Temperature (°C)  
Std Dev < Dotted >  
February  
1000 Mb



Mean Temperature (°C)



Std Dev < Dotted >

February

850 Mb

Upper Air Climatology

Northern Hemisphere



Upper Air Climatology



Southern Hemisphere

Mean Temperature (°C)

Std Dev < Dotted >

February

850 Mb



Mean Temperature (°C)



Std Dev < Dotted >

February

700 Mb



### Upper Air Climatology

Northern Hemisphere



Upper Air Climatology  
Southern Hemisphere

Mean Temperature (c)  
Std Dev <Dotted>  
February  
700 Mb



Mean Temperature (°C)


Std Dev <Dotted>

February

500 Mb



Upper Air Climatology

Northern Hemisphere



Upper Air Climatology  
Southern Hemisphere

Mean Temperature (c)  
Std Dev <Dotted>  
February  
500 Mb





Mean Temperature (c)

Std Dev < Dotted >

February


400 Mb

Upper Air Climatology  
Northern Hemisphere



Upper Air Climatology  
Southern Hemisphere

Mean Temperature (c)  
Std Dev < Dotted >  
February  
400 Mb




### Mean Temperature (c)

Std Dev <Dotted>

February


300 Мъ

## Upper Air Climatology Northern Hemisphere



Upper Air Climatology  
Southern Hemisphere

Mean Temperature (c)  
Std Dev < Dotted >  
February  
300 MB



Mean Temperature (°C)



Std Dev <Dotted>

February

250 Mb


Upper Air Climatology

Northern Hemisphere



Upper Air Climatology  
Southern Hemisphere

Mean Temperature (°C)  
Std Dev < Dotted>  
February  
250 Mb



Mean Temperature (°C)


Std Dev < Dotted >

February

200 MB


Upper Air Climatology

Northern Hemisphere




Upper Air Climatology  
Southern Hemisphere

Mean Temperature (°C)  
Std Dev < Dotted >  
February  
200 Mb



Std. Dev. < 2.5



Mean Temperature (°C)



Std Dev <Dotted>

February

150 Mb


Upper Air Climatology

Northern Hemisphere




Upper Air Climatology  
Southern Hemisphere

Mean Temperature (c)  
Std Dev < Dotted >  
February  
150 Mb



Std. Dev. < 2.5



Std. Dev. < 2.5



Mean Temperature (c)

Std Dev < Dotted >

February



100 Mb

Upper Air Climatology  
Northern Hemisphere



Upper Air Climatology  
Southern Hemisphere

Mean Temperature (°C)  
Std Dev < Dotted  
February  
100 Mb



Mean Temperature (°C)



Std Dev <Dotted>

February

70 Mb



Upper Air Climatology

Northern Hemisphere



Upper Air Climatology  
Southern Hemisphere

Mean Temperature (°C)  
Std Dev < Dotted >  
February  
70 Mb



Mean Temperature (c)



Std Dev < Dotted >

February

50 Mb

Upper Air Climatology

Northern Hemisphere



Upper Air Climatology


Southern Hemisphere

Mean Temperature (c)

Std Dev < Dotted >

February

50 Mb



Mean Temperature (c)


Std Dev < Dotted >

February

30 Mb


Upper Air Climatology

Northern Hemisphere




Upper Air Climatology  
Southern Hemisphere

Mean Temperature (°C)  
Std Dev < Dotted >  
February  
30 Mb




Std. Dev. > 2.5



**DEW POINT**  
**(6 LEVELS, 1000 TO 300 MB)**

- Contours of mean dew point (solid and dashed lines) in °C; solids labeled, dashed intermediates unlabeled.
- Dew point labeled interval: 5°C
- Contours of standard deviation of dew point (dotted lines) in °C
- Standard deviation of dew point labeled interval: 2.5°C
- Contours blanked for geographic areas with elevations exceeding specified geopotential heights

**ELEVATION SCALE**



Mean Dew Point (°C)


Std Dev (Dotted)

February

1000 MB

Upper Air Climatology

Northern Hemisphere



Upper Air Climatology



Southern Hemisphere

Mean Dew Point (c)

Std Dev <Dotted>

February

1000 Mb



### Mean Dew Point (c)

### Std Dev <Dotted>

February

350 MB

## Upper Air Climatology Northern Hemisphere





Upper Air Climatology  
Southern Hemisphere

Mean Dew Point (c)

Std Dev < Dotted >

February

850 MB



Mean Dew Point (c)



Std Dev < Dotted >

February

700 Mb



Upper Air Climatology

Northern Hemisphere



Upper Air Climatology  
Southern Hemisphere

Mean Dew Point (c)  
Std Dev < Dotted >  
February  
700 Mb





Mean Dew Point (c)

Std Dev < Dotted >

February

500 Mb

Upper Air Climatology  
Northern Hemisphere





Upper Air Climatology  
Southern Hemisphere

Mean Dew Point (c)

Std Dev < Dotted >

February

500 Mb



Mean Dew Point (c)



Std Dev < Dotted >

February

400 MB


Upper Air Climatology

Northern Hemisphere



Upper Air Climatology  
Southern Hemisphere

Mean Dew Point (°C)  
Std Dev < Dotted >  
February  
400 Mb



Mean Dew Point (c)



Std Dev < Dotted >

February

300 Mb


Upper Air Climatology

Northern Hemisphere



Upper Air Climatology  
Southern Hemisphere


Mean Dew Point (c)  
Std Dev < Dotted>  
February  
300 Mb



**DENSITY**  
**(13 LEVELS, 1000 TO 30 MB)**

- Contours of mean density (solid and dashed lines) in kilograms/cubic meter; solids labeled, dashed intermediates unlabeled
- Density labeled interval:
  - .02 kilograms/cubic meter - 1000 MB to 400 MB
  - .01 kilograms/cubic meter - 300 MB to 200 MB
  - .006 kilograms/cubic meter - 150 MB to 30 MB
- Contours of standard deviation of density (dotted lines) in kilograms/cubic meter
- Standard deviation of density labeled interval:
  - .01 kilograms/cubic meter - 1000 MB to 400 MB
  - .005 kilograms/cubic meter - 300 MB to 200 MB
  - .003 kilograms/cubic meter - 150 MB to 30 MB
- Contours blanked for geographic areas with elevations exceeding specified geopotential heights

**ELEVATION SCALE**



Mean Density ( $\text{kg/m}^3$ )


Std Dev <Dotted>

February

1000 Mb



Upper Air Climatology

Northern Hemisphere



Upper Air Climatology  
Southern Hemisphere

Mean Density ( $\text{kg/m}^3$ )  
Std Dev <Dotted>  
February  
1000 Mb



Mean Density ( $\text{kg/m}^3$ )


Std Dev < Dotted >

February

850 Mb


Upper Air Climatology

Northern Hemisphere



Upper Air Climatology  
Southern Hemisphere

Mean Density ( $\text{kg/m}^3$ )  
Std Dev <Dotted>  
February  
850 Mb



Mean Density ( $\text{kg/m}^3$ )


Std Dev <Dotted>

February

700 MB


Upper Air Climatology

Northern Hemisphere



Upper Air Climatology  
Southern Hemisphere

Mean Density ( $\text{kg/m}^3$ )  
Std Dev < Dotted >  
February  
700 Mb



### Mean Density (kg/m<sup>3</sup>)


### Std Dev <Dotted>

February

500 Mb



## Upper Air Climatology

### Northern Hemisphere



Upper Air Climatology  
Southern Hemisphere

Mean Density ( $\text{kg/m}^3$ )  
Std Dev < Dotted >  
February  
500 Mb



Mean Density ( $\text{kg/m}^3$ )



Std Dev < Dotted >

February

400 MB


Upper Air Climatology

Northern Hemisphere



Upper Air Climatology  
Southern Hemisphere

Mean Density ( $\text{kg/m}^3$ )  
Std Dev < Dotted >  
February  
400 Mb



Mean Density ( $\text{kg/m}^3$ )


Std Dev < Dotted >

February

300 Mb


Upper Air Climatology

Northern Hemisphere



Upper Air Climatology  
Southern Hemisphere

Mean Density ( $\text{kg/m}^3$ )  
Std Dev < Dotted  
February  
300 Mb



Mean Density ( $\text{kg/m}^3$ )



Std Dev < Dotted >

February

250 Mb

Upper Air Climatology

Northern Hemisphere




## Upper Air Climatology Southern Hemisphere

### Mean Density (kg/m<sup>3</sup>)

Std Dev <Dotterá>

February

250 MB



Mean Density ( $\text{kg/m}^3$ )



Std Dev <Dotted>

February

200 Mb



Upper Air Climatology

Northern Hemisphere



Upper Air Climatology  
Southern Hemisphere

Mean Density ( $\text{kg/m}^3$ )  
Std Dev < Dotted  
February  
200 Mb



Mean Density ( $\text{kg/m}^3$ )



Std Dev < Dotted >

February

150 Mb

Upper Air Climatology

Northern Hemisphere



Upper Air Climatology



Southern Hemisphere

Mean Density ( $\text{kg/m}^3$ )

Std Dev < Dotted>

February

150 Mb



Mean Density ( $\text{kg/m}^3$ )



Std Dev < Dotted >

February

100 Mb

Upper Air Climatology

Northern Hemisphere





Upper Air Climatology  
Southern Hemisphere

Mean Density ( $\text{kg/m}^3$ )

Std Dev < Dotted >

February

100 Mb





### Mean Density (kg/m<sup>3</sup>)

### **Std Dev <Dotted>**

February



70 Mb

## Upper Air Climatology Northern Hemisphere



## Upper Air Climatology Southern Hemisphere

Mean Density (kg/m<sup>3</sup>)  
Std Dev < Dotted >  
February  
70 Mb



Mean Density ( $\text{kg/m}^3$ )



Std Dev < Dotted >

February

50 Mb



Upper Air Climatology

Northern Hemisphere



Upper Air Climatology  
Southern Hemisphere

Mean Density ( $\text{kg/m}^3$ )  
Std Dev < Dotted >  
February  
50 Mb



Mean Density ( $\text{kg/m}^3$ )



Std Dev < Dotted >

February

30 Mb


Upper Air Climatology

Northern Hemisphere



Upper Air Climatology  
Southern Hemisphere


Mean Density ( $\text{kg}/\text{m}^3$ )  
Std Dev < Dotted>  
February  
30 Mb



**STANDARD DEVIATION OF HEIGHT  
STANDARD DEVIATION OF VECTOR MEAN WIND  
(13 LEVELS, 1000 TO 30 MB)**

- Contours of standard deviation of height (solid lines) in geopotential dekameters
- Standard deviation of height labeled interval:
  - 3 dekameters (30 meters) - 1000 MB to 400 MB
  - 6 dekameters (60 meters) - 300 MB to 200 MB
  - 4 dekameters (40 meters) - 150 MB to 30 MB
- Contours of standard deviation of vector mean wind (dashed lines) in knots
- Standard deviation of vector mean wind labeled interval: 5 knots
- Contours blanked for geographic areas with elevations exceeding specified geopotential heights

**ELEVATION SCALE**



Height (dkm) Std Dev <Solid>


Vector Std Dev (kt)

February

1000 Mb


Upper Air Climatology

Northern Hemisphere



Upper Air Climatology  
Southern Hemisphere

Height (dkm) Std Dev <Solid>  
Vector Std. Dev (kt)  
February  
1000 Mb



Height (dkm) Std Dev <Solid>



Vector Std Dev (kt)

February

850 Mb

Upper Air Climatology

Northern Hemisphere




Upper Air Climatology  
Southern Hemisphere

Height (dkm) Std Dev <Solid>

Vector Std Dev (kt)

February

850 Mb



Height (dkm) Std Dev <Solid>



Vector Std Dev (kt)

February

700 MB



Upper Air Climatology

Northern Hemisphere



Upper Air Climatology  
Southern Hemisphere

Height (dkm) Std Dev <Solid>  
Vector Std Dev (kt)  
February  
700 Mb



Height (dkm) Std Dev <Solid>


Vector Std Dev (kt)

February

500 Mb



Upper Air Climatology

Northern Hemisphere



Upper Air Climatology  
Southern Hemisphere

Height (dkm) Std Dev <Solid>  
Vector Std Dev (kt)  
February  
500 MB



Height (dkm) Std Dev <Solid>



Vector Std Dev (kt)

February

400 Mb

Upper Air Climatology

Northern Hemisphere



Upper Air Climatology


Southern Hemisphere

Height (dkm) Std Dev <Solid>

Vector Std Dev (kt)

February

400 Mb




Height (dkm) Std Dev <Solid>

Vector Std Dev (kt)

February

300 MB

Upper Air Climatology  
Northern Hemisphere





Upper Air Climatology  
Southern Hemisphere

Height (dkm) Std Dev <Solid>

Vector Std Dev (kt)

February

300 Mb



Height (dkm) Std Dev <Solid>



Vector Std Dev (kt)

February

250 Mb

Upper Air Climatology

Northern Hemisphere



Upper Air Climatology


Southern Hemisphere

Height (dkm) Std Dev <Solid>

Vector Std Dev (kt)

February

250 Mb



Height (dkm) Std Dev <Solid>


Vector Std Dev (kt)

February

210 MS

Upper Air Climatology

Northern Hemisphere



Upper Air Climatology



Southern Hemisphere

Height (dkm) Std Dev <Solid>

Vector Std Dev (kt)

February

200 Mb



Height (dkm) Std Dev < Solid>



Vector Std Dev (kt)

February

850 Mb



### Upper Air Climatology

#### Northern Hemisphere



Upper Air Climatology  
Southern Hemisphere

Height (dkm) Std Dev < Solid>  
Vector Std Dev (kt)  
February  
150 Mb



Height (dkm) Std Dev <Solid>



Vector Std Dev (kt)

February

100 Mb

Upper Air Climatology

Northern Hemisphere




Upper Air Climatology  
Southern Hemisphere

Height (dkm) Std Dev < Solid>

Vector Std Dev (kt)

February

100 Mb



Height (dkm) Std Dev <Solid>


## Upper Air Climatology

### Northern Hemisphere

### Vector Std Dev (kt)

February

70 Mb



Upper Air Climatology


Southern Hemisphere

Height (dkm) Std Dev < Solid>

Vector Std Dev (kt)

February

70 Mb



Height Std. Dev. < 12


Vector Std. Dev. < 10.0

10.0

10.0 10.0

10.0 Height Std. Dev. < 12

Vector Std. Dev. < 10.0



Height (dkm) Std Dev <Solid>


Vector Std Dev (kt)

February

50 Mb

Upper Air Climatology

Northern Hemisphere




Upper Air Climatology  
Southern Hemisphere

Height (dkm) Std Dev < Solid>


Vector Std Dev (kt)

February

50 Mb



Height Std. Dev. < 12



Height (dkm) Std Dev <Solid>


Vector Std Dev (kt)

February

20 MB

Upper Air Climatology

Northern Hemisphere



Upper Air Climatology  
Southern Hemisphere

Height (dkm) Std Dev <Solid>  
Vector Std Dev (kt)  
February  
30 Mb

